

`$SPAD/input schaum3.input`

Timothy Daly

June 15, 2008

## Contents

<b>1</b>	<b>[1]:14.105</b>	$\int \frac{dx}{(ax+b)(px+q)}$	<b>3</b>
<b>2</b>	<b>[1]:14.106</b>	$\int \frac{x \ dx}{(ax+b)(px+q)}$	<b>5</b>
<b>3</b>	<b>[1]:14.107</b>	$\int \frac{dx}{(ax+b)^2(px+q)}$	<b>6</b>
<b>4</b>	<b>[1]:14.108</b>	$\int \frac{x \ dx}{(ax+b)^2(px+q)}$	<b>8</b>
<b>5</b>	<b>[1]:14.109</b>	$\int \frac{x^2 \ dx}{(ax+b)^2(px+q)}$	<b>10</b>
<b>6</b>	<b>[1]:14.110</b>	$\int \frac{dx}{(ax+b)^m(px+q)^n}$	<b>11</b>
<b>7</b>	<b>[1]:14.111</b>	$\int \frac{ax+b}{px+q} \ dx$	<b>12</b>
<b>8</b>	<b>[1]:14.112</b>	$\int \frac{(ax+b)^m}{(px+q)^n} \ dx$	<b>13</b>

**1 [1]:14.105**       $\int \frac{dx}{(ax+b)(px+q)}$   

$$\int \frac{1}{(ax+b)(px+q)} = \frac{1}{bp-aq} \ln \left( \frac{px+q}{ax+b} \right)$$

```

(*)≡
)spool schaum3.output
)set message test on
)set message auto off
)clear all

--S 1
aa:=integrate(1/((a*x+b)*(p*x+q)),x)
--R
--R
--R      - log(p x + q) + log(a x + b)
--R      (1)  -----
--R                  a q - b p
--R
--E                                         Type: Union(Expression Integer,...)

--S 2
bb:=1/(b*p-a*q)*log((p*x+q)/(a*x+b))
--R
--R
--R      p x + q
--R      log(-----)
--R                  a x + b
--R      (2)  -
--R                  a q - b p
--R
--E                                         Type: Expression Integer

--S 3
cc:=aa-bb
--R
--R
--R      p x + q
--R      - log(p x + q) + log(a x + b) + log(-----)
--R                                              a x + b
--R      (3)  -----
--R                  a q - b p
--R
--E                                         Type: Expression Integer

```

```

--S 4
logdiv:=rule(log(a)-log(b) == log(a/b))
--R
--R
--R      (4)  - log(b) + log(a) + %I == log(-) + %I
--R                                a
--R                                b
--R                                         Type: RewriteRule(Integer,Integer,Expression Integer)
--E

--S 5
dd:=logdiv cc
--R
--R
--R      (5)  -----
--R          log(a x + b) + log(-----)
--R                      a x + b
--R          a q - b p
--R                                         Type: Expression Integer
--E

--S 6
logmul:=rule(log(a)+log(b) == log(a*b))
--R
--R      (6)  log(b) + log(a) + %J == log(a b) + %J
--R                                         Type: RewriteRule(Integer,Integer,Expression Integer)
--E

--S 7      14:105 Schaums and Axiom agree
ee:=logmul dd
--R
--R      (7)  0
--R                                         Type: Expression Integer
--E

```

**2 [1]:14.106**       $\int \frac{x \, dx}{(ax+b)(px+q)}$   

$$\int \frac{x}{(ax+b)(px+q)} = \frac{1}{bp-aq} \left\{ \frac{b}{a} \ln(ax+b) - \frac{q}{p} \ln(px+q) \right\}$$

$$\langle * \rangle + \equiv$$

$$)\text{clear all}$$
  
--S 8
aa:=integrate(x/((a\*x+b)\*(p\*x+q)),x)
--R
--R
--R (1) 
$$\frac{a q \log(p x + q) - b p \log(a x + b)}{a^2 p^2 q^2 - a^2 b^2 p^2}$$
--R
--R
--E
 Type: Union(Expression Integer,...)
  
--S 9
bb:=1/(b\*p-a\*q)\*(b/a\*log(a\*x+b)-q/p\*log(p\*x+q))
--R
--R
--R (2) 
$$\frac{a q \log(p x + q) - b p \log(a x + b)}{a^2 p^2 q^2 - a^2 b^2 p^2}$$
--R
--R
--E
 Type: Expression Integer
  
--S 10 14:106 Schaums and Axiom agree
cc:=aa-bb
--R
--R
--R (3) 0
--R
--E
 Type: Expression Integer

**3 [1]:14.107**

$$\int \frac{dx}{(ax+b)^2(px+q)}$$

$$\int \frac{1}{(ax+b)^2(px+q)} = \frac{1}{bp-aq} \left\{ \frac{1}{ax+b} + \frac{p}{bp-aq} \ln \left( \frac{px+q}{ax+b} \right) \right\}$$

(\*)+≡  
 )clear all

--S 11  
 aa:=integrate(1/((a\*x+b)^2\*(p\*x+q)),x)  
 --R  
 --R  
 --R (1) 
$$\frac{(a p x + b p) \log(p x + q) + (- a p x - b p) \log(a x + b) - a q + b p}{(a q^3 - 2 a b p q^2 + a b p^2)x^3 + a b q^2 - 2 a b p q + b p^2}$$
  
 Type: Union(Expression Integer,...)  
 --E

--S 12  
 bb:=1/(b\*p-a\*q)\*(1/(a\*x+b)+p/(b\*p-a\*q)\*log((p\*x+q)/(a\*x+b)))  
 --R  
 --R  
 --R (2) 
$$\frac{(a p x + b p) \log(\frac{p x + q}{a x + b}) - a q + b p}{(a q^3 - 2 a b p q^2 + a b p^2)x^3 + a b q^2 - 2 a b p q + b p^2}$$
  
 Type: Expression Integer  
 --E

--S 13  
 cc:=aa-bb  
 --R  
 --R  
 --R (3) 
$$\frac{p \log(p x + q) - p \log(a x + b) - p \log(\frac{p x + q}{a x + b})}{(a q^2 - 2 a b p q + b p^2)}$$
  
 Type: Expression Integer  
 --E

```

--S 14
divlog:=rule(log(a/b) == log(a) - log(b))
--R
--R          a
--R      (4)  log(-) == - log(b) + log(a)
--R          b
--R                                         Type: RewriteRule(Integer,Integer,Expression Integer)
--E

--S 15      14:107 Schaums and Axiom agree
dd:=divlog cc
--R
--R      (5)  0
--R                                         Type: Expression Integer
--E

```

4 [1]:14.108

$$\int \frac{x \, dx}{(ax+b)^2(px+q)}$$

$$\int \frac{x}{(ax+b)^2(px+q)} = \frac{1}{bp-aq} \left\{ \frac{q}{bp-aq} \ln \left( \frac{ax+b}{px+q} \right) - \frac{b}{a(ax+b)} \right\}$$

$\langle *\rangle + \equiv$

)clear all

--S 16

```
aa:=integrate(x/((a*x+b)^2*(p*x+q)),x)
--R
--R
--R      (1)
```

$$\frac{(-a q x^2 - a b q) \log(p x^2 + q) + (a q x^3 + a b q) \log(a x^2 + b) + a b q^2 - b p^2}{(a q^4 - 2 a b p q^3 + a b p^2)x^4 + a b q^3 - 2 a b p q^2 + a b p^2}$$

Type: Union(Expression Integer,...)

--E

--S 17

```
bb:=1/(b*p-a*q)*(q/(b*p-a*q)*log((a*x+b)/(p*x+q))-b/(a*(a*x+b)))
--R
--R
--R      (2)
```

$$\frac{(a q x^2 + a b q) \log(\frac{a x^2 + b}{p x^2 + q})^2 + a b q^2 - b p^2}{(a q^4 - 2 a b p q^3 + a b p^2)x^4 + a b q^3 - 2 a b p q^2 + a b p^2}$$

Type: Expression Integer

--E

--S 18

```
cc:=aa-bb
--R
--R
--R      (3)
```

$$\frac{-q \log(p x^2 + q) + q \log(a x^2 + b) - q \log(\frac{a x^2 + b}{p x^2 + q})}{(a q^2 - 2 a b p q + b p^2)^2}$$

Type: Expression Integer

```

--E

--S 19
divlog:=rule(log(a/b) == log(a) - log(b))
--R
--R          a
--R      (4)  log(--) == - log(b) + log(a)
--R          b
--R                                         Type: RewriteRule(Integer,Integer,Expression Integer)
--E

--S 20      14:108 Schaums and Axiom agree
dd:=divlog cc
--R
--R      (5)  0
--R                                         Type: Expression Integer
--E

```

5 [1]:14.109

$$\int \frac{x^2}{(ax+b)^2(px+q)} dx$$

$$\int \frac{x^2}{(ax+b)^2(px+q)} =$$

$$\frac{b^2}{(bp-aq)a^2(ax+b)} + \frac{1}{(bp-aq)^2} \left\{ \frac{q^2}{p} \ln(px+q) + \frac{b(bp-2aq)}{a^2} \ln(ax+b) \right\}$$

$\langle *\rangle + \equiv$

)clear all

--S 21

```
aa:=integrate(x^2/((a*x+b)^2*(p*x+q)),x)
--R
--R
--R (1)
```

$$(a q x + a b q) \log(p x + q)$$

$$+ \frac{((-2a b p q + a b p)x^2 - 2a b p q + b p) \log(a x + b) - a b p q + b p}{(a p q^5 - 2a b p q^4 + a b p^2)x^3 + a b p q^4 - 2a b p q^3 + a b p^2}$$

Type: Union(Expression Integer,...)

--E

--S 22

```
bb:=b^2/((b*p-a*q)*a^2*(a*x+b))+_
1/(b*p-a*q)^2*(q^2/p*log(p*x+q)+((b*(b*p-2*a*q))/a^2)*log(a*x+b))
--R
--R
--R (2)
```

$$(a q x + a b q) \log(p x + q)$$

$$+ \frac{((-2a b p q + a b p)x^2 - 2a b p q + b p) \log(a x + b) - a b p q + b p}{(a p q^5 - 2a b p q^4 + a b p^2)x^3 + a b p q^4 - 2a b p q^3 + a b p^2}$$

Type: Expression Integer

--E

--S 23 14:109 Schaums and Axiom agree

```

cc:=aa-bb
--R
--R
--R      (3)  0
--R
--E                                         Type: Expression Integer

6   [1]:14.110      
$$\int \frac{dx}{(ax+b)^m(px+q)^n}$$


$$\int \frac{1}{(ax+b)^m(px+q)^n} =$$


$$\frac{-1}{(n-1)(bp-aq)} \left\{ \frac{1}{(ax+b)^{m-1}(px+q)^{n-1}} + a(m+n-2) \int \frac{1}{(ax+b)^m(px+q)^{n-1}} \right\}$$


$$\langle * \rangle + \equiv$$

)clear all

--S 24      14:110 Axiom cannot do this integral
aa:=integrate(1/((a*x+b)^m*(p*x+q)^n),x)
--R
--R
--R      x
--R      ++
--R      (1)  |  -----
--R      ++          m           n
--R                  (b + %L a) (q + %L p)                                         Type: Union(Expression Integer,...)
--E

```

**7 [1]:14.111**  $\int \frac{ax+b}{px+q} dx$   

$$\int \frac{ax+b}{px+q} = \frac{ax}{p} + \frac{bp-aq}{p^2} \ln(px+q)$$

```

(*)+≡
)clear all

--S 25
aa:=integrate((a*x+b)/(p*x+q),x)
--R
--R
--R      (- a q + b p)log(p x + q) + a p x
--R      (1)  -----
--R                  2
--R                  p
--R                                         Type: Union(Expression Integer,...)
--E

--S 26
bb:=(a*x)/p+(b*p-a*q)/p^2*log(p*x+q)
--R
--R
--R      (- a q + b p)log(p x + q) + a p x
--R      (2)  -----
--R                  2
--R                  p
--R                                         Type: Expression Integer
--E

--S 27      14:111 Schaums and Axiom agree
cc:=aa-bb
--R
--R
--R      (3)  0
--R                                         Type: Expression Integer
--E

```

8 [1]:14.112

$$\int \frac{(ax+b)^m}{(px+q)^n} dx$$

$$\int \frac{(ax+b)^m}{(px+q)^n} = \begin{cases} \frac{-1}{(n-1)(bp-aq)} \left\{ \frac{(ax+b)^{m+1}}{(px+q)^{n-1}} + (n-m-2)a \int \frac{(ax+b)^m}{(px+q)^{n-1}} \right\} \\ \frac{-1}{(n-m-1)p} + \left\{ \frac{(ax+b)^m}{(px+q)^{n-1}} + m(bp-aq) \int \frac{(ax+b)^{m-1}}{(px+q)^n} \right\} \\ \frac{-1}{(n-1)p} \left\{ \frac{(ax+b)^m}{(px+q)^{n-1}} - ma \int \frac{(ax+b)^{m-1}}{(px+q)^{n-1}} \right\} \end{cases}$$

$\langle *\rangle + \equiv$

)clear all

--S 28 14:112 Axiom cannot do this integral

aa:=integrate((a\*x+b)^m/(p\*x+q)^n,x)

--R

--R

--R

--I                x                m

--I                ++ (b + %L a)               

--I        (1)     | ----- d%L

--R                ++                n

--I                (q + %L p)

--R

Type: Union(Expression Integer,...)

--E

$\langle *\rangle + \equiv$

)spool

)lisp (bye)

## References

- [1] Spiegel, Murray R. *Mathematical Handbook of Formulas and Tables*  
Schaum's Outline Series McGraw-Hill 1968 pp62-63